

Hall operators on the set of formations of finite groups

Andrei P. Mekhovich, Nikolay N. Vorob'ev
and Nikolay T. Vorob'ev

Communicated by L. A. Shemetkov

*Dedicated to Professor I. Ya. Subbotin
on the occasion of his 60-th birthday*

ABSTRACT. Let π be a nonempty set of primes and let \mathfrak{F} be a saturated formation of all finite soluble π -groups. It is constructed the saturated formation consisting of all finite π -soluble groups whose \mathfrak{F} -projectors contain a Hall π -subgroup.

Introduction

In the theory of soluble Fitting classes P. Lockett and P. Hauck considered the classes $\mathcal{L}_\pi(\mathfrak{F})$ and $\mathcal{K}_\pi(\mathfrak{F})$.

Definition 1 ([1, 2]). *Let π be a set of primes and let \mathfrak{F} be a Fitting class of finite soluble groups. Then*

$$\begin{aligned}\mathcal{L}_\pi(\mathfrak{F}) &= (G \in \mathfrak{S} : \text{an } \mathfrak{F}\text{-injector of } G \text{ contains a Hall } \pi\text{-subgroup of } G); \\ \mathcal{K}_\pi(\mathfrak{F}) &= (G \in \mathfrak{S} : \text{a Hall } \pi\text{-subgroup of } G \text{ belongs to } \mathfrak{F}).\end{aligned}$$

In [1] (see also [3, IX, 1.22]) Lockett used the class $\mathcal{L}_\pi(\mathfrak{F})$ to obtain a description of the injectors for a Fitting class product $\mathfrak{F}\mathfrak{G}$. It was proved that $\mathcal{L}_\pi(\mathfrak{F})$ and $\mathcal{K}_\pi(\mathfrak{F})$ are Fitting classes. Furthermore, $\mathcal{K}_\pi(\mathfrak{F}) =$

Research of the second author is partially supported by Belarussian Republic Foundation of Fundamental Researches (BRFFI, grant F08M-118)

2001 Mathematics Subject Classification: 20D10.

Key words and phrases: Hall π -subgroup, π -soluble group, formation of finite groups, saturated formation, canonical satellite, \mathfrak{F} -projector.

$\mathcal{L}_\pi(\mathfrak{F} \cap \mathfrak{S}_\pi)$. Hence we may consider \mathcal{L}_π and \mathcal{K}_π as operators on the set of all Fitting classes for every π . The class $\mathcal{K}_\pi(\mathfrak{F})$ was introduced by Hauck [2] and has been studied in detail by Brison [4] and Cusack [5]. Moreover, Brison [6, 7] applied $\mathcal{K}_\pi(\mathfrak{F})$ to obtain a description of Hall subgroups radicals.

Analogously one may consider the following operators on the set of all soluble formations.

Definition 2 ([8, 9]). *Let π be a set of primes and let \mathfrak{F} be a formation of finite soluble groups. Then*

$$\mathcal{L}^\pi(\mathfrak{F}) = (G \in \mathfrak{S} : \text{an } \mathfrak{F}\text{-projector of } G \text{ contains a Hall } \pi\text{-subgroup of } G);$$

$$\mathcal{K}^\pi(\mathfrak{F}) = (G \in \mathfrak{S} : \text{a Hall } \pi\text{-subgroup of } G \text{ belongs to } \mathfrak{F}).$$

In [9] Blessenohl proved that if \mathfrak{F} is a saturated formation, then $\mathcal{K}^\pi(\mathfrak{F})$ is a saturated formation.

Further L.A. Shemetkov posed the following question in this trend.

Problem (see [10, Problem 19]). Let \mathfrak{F} be a saturated formation of finite groups, $\mathcal{C}_\pi(\mathfrak{F})$ be the class of all groups G such that there exist Hall π -subgroups of G in \mathfrak{F} and any two of them are conjugate. Is the class $\mathcal{C}_\pi(\mathfrak{F})$ a saturated formation?

The positive answer of Problem 19 was given by L.M. Slepova [11] in the class of all π -separable groups for some restrictions to \mathfrak{F} ; in [12] it was shown by E.P. Vdovin, D.O. Revin and L.A. Shemetkov that $\mathcal{C}_\pi(\mathfrak{F})$ is solubly saturated formation for any solubly saturated formation \mathfrak{F} . However L.A. Shemetkov and A.F. Vasil'ev [13] proved that in general the class $\mathcal{C}_\pi(\mathfrak{N})$ is not a saturated formation, where \mathfrak{N} is the class of all nilpotent groups.

Wenbin Guo and Baojin Li [14] proved that $\mathcal{K}_\pi(\mathfrak{F})$ is a local Fitting class for every local Fitting class \mathfrak{F} . In general N.T. Vorob'ev and V.N. Zagurskii [15] gave the positive answer of Shemetkov's Problem for soluble ω -local Fitting classes.

K. Doerk and T. Hawkes investigated an analog of Problem 19 for the class $\mathcal{L}^\pi(\mathfrak{F})$. It was proved, that if \mathfrak{F} is a solubly saturated formation, then $\mathcal{L}^\pi(\mathfrak{F})$ is a saturated formation (see [8, Bemerkung]). Note that the analog of the above-mentioned problem has the negative answer for soluble Schunck classes (see [8, Beispiel 1]) and soluble Fitting classes (see [3, IX, 3.15]).

A purpose of this paper is to investigate an analog of Shemetkov's Problem for the class $\mathcal{L}^\pi(\mathfrak{F})$, where \mathfrak{F} is the saturated formation of all soluble π -groups.

All groups considered are finite and π -soluble for some fixed nonempty set of primes π . All unexplained notations and terminologies are standard. The reader is referred to [16], [10] and [3] if necessary.

1. Preliminaries

Recall notation and some definitions used in this paper.

A group class closed under taking homomorphic images and finite subdirect products is called a *formation*.

A group G is said to be π -soluble if every chief factor of G is either a p -group for some $p \in \pi$ or a π' -subgroup.

The complementary set of primes, $\mathbb{P} \setminus \pi$, is denoted by π' . $\sigma(G)$ denotes the set of all distinct prime divisors of the order of a group G .

Functions of the form

$$f : \mathbb{P} \rightarrow \{\text{formations of groups}\}$$

are called *local satellites* (see [10]). For every local satellite f it is defined the class

$$LF(f) = (G : G \text{ has } f\text{-central chief series}),$$

i.e., for every chief factor H/K of G we have

$$G/C_G(H/K) \in f(p) \text{ for every } p \in \pi(H/K).$$

If \mathfrak{F} is a formation such that $\mathfrak{F} = LF(f)$ for a local satellite f , then the formation \mathfrak{F} is said to be *saturated* and f is a local satellite of \mathfrak{F} .

If \mathfrak{F} is a saturated formation, by [3, IV, 4.3] we have $\text{Char}(\mathfrak{F}) = \sigma(\mathfrak{F})$, where $\sigma(\mathfrak{F}) = \bigcup \{\sigma(G) : G \in \mathfrak{F}\}$.

A satellite F of a formation \mathfrak{F} is called *canonical* if $F(p) \subseteq \mathfrak{F}$, and $F(p) = \mathfrak{N}_p F(p)$ for all $p \in \mathbb{P}$ [17].

Let \mathfrak{F} be a formation. A subgroup H of a group G is called \mathfrak{F} -maximal in G provided that

- (1) $H \in \mathfrak{F}$, and
- (2) if $H \leq V \leq G$ and $V \in \mathfrak{F}$, then $H = V$.

A subgroup H of G is called an \mathfrak{F} -projector of G if HN/N is \mathfrak{F} -maximal in G/N for all $N \trianglelefteq G$.

By $\text{Proj}_{\mathfrak{F}}G$ we denote the (possibly empty) set of all \mathfrak{F} -projectors of G .

Let \mathfrak{F} be a saturated formation and let \mathfrak{H} be a formation. Following [3, IV, 1.1] we denote the class $(\mathfrak{F} \downarrow \mathfrak{H})$ as follows:

$$(\mathfrak{F} \downarrow \mathfrak{H}) = (G : \text{Proj}_{\mathfrak{F}}G \subseteq \mathfrak{H}).$$

If $\mathfrak{H} = \emptyset$, then $(\mathfrak{F} \downarrow \mathfrak{H}) = \emptyset$.

If $RB \supseteq A$, then it is said that A/B covered by R .

The symbols G_π , \mathfrak{S}^π , $\mathfrak{E}_{\pi'}$, \mathfrak{E}_π and \mathfrak{N}_p denote, respectively, a Hall π -subgroup of a group G , the class of all π -soluble groups, the class of all π' -groups, the class of all π -groups and the class of all p -groups.

We need some lemmas to prove the main result.

Lemma 1 ([18, Lemma 1.2, Lemma 1.3]). *Let $\mathfrak{F} = LF(F)$ be the formation of all soluble π -groups. Then the following statements hold:*

(1) $\mathfrak{F} = LF(m)$, where

$$m(p) = (\mathfrak{F} \downarrow F(p)) \text{ for all } p \in \mathbb{P}.$$

(2) *If V is an \mathfrak{F} -projector of a group G , then:*

(a) V covers every m -central chief factor of G .

(b) Every chief factor of G covered of the subgroup V is m -central.

Lemma 2 ([10, Theorem 15.7]). *Let \mathfrak{F} be a saturated formation and G be a group having $\sigma(\mathfrak{F})$ -soluble \mathfrak{F} -residual. Then G has \mathfrak{F} -projectors and any two of them are conjugate.*

2. The proof of Theorem

First we prove

Lemma 3. *Let \mathfrak{F} be a saturated formation of all soluble π -groups. Then the following statements hold:*

(1) *The class $\mathcal{L}^\pi(\mathfrak{F})$ is a formation.*

(2) $\mathfrak{E}_{\pi'} \mathcal{L}^\pi(\mathfrak{F}) = \mathcal{L}^\pi(\mathfrak{F})$.

Proof. (1) If $\pi = \emptyset$, then $\mathcal{L}^\emptyset(\mathfrak{F}) = \mathfrak{S}^\pi$; if $\pi = \mathbb{P}$, then $\mathcal{L}^\mathbb{P}(\mathfrak{F}) = \mathfrak{F}$. We have saturated formations \mathfrak{S}^π and \mathfrak{F} , and hence the result. Now suppose $\emptyset \subset \pi \subset \mathbb{P}$. Since a formation \mathfrak{F} is saturated, by [3, IV, 4.3] we have $\text{Char}(\mathfrak{F}) = \sigma(\mathfrak{F})$.

Since $\sigma(\mathfrak{F}) \subseteq \pi$, a π -soluble group G is $\sigma(\mathfrak{F})$ -soluble. Consequently, the subgroup $G^{\mathfrak{F}}$ of G is $\sigma(\mathfrak{F})$ -soluble.

Let $G \in \mathcal{L}^\pi(\mathfrak{F})$, let $K \triangleleft G$ and let F be an \mathfrak{F} -projector of G . Then there exists a Hall π -subgroup G_π of G such that $G_\pi \subseteq F$.

By [10, Lemma 15.2] and [10, Lemma 15.1], we see that $G_\pi K/K$ is a Hall π -subgroup of G/K and FK/K is an \mathfrak{F} -projector of G/K . Therefore $G/K \in \mathcal{L}^\pi(\mathfrak{F})$.

Let K_1 and K_2 be normal subgroups of G such that $K_1 \cap K_2 = 1$ and let $G/K_1 \in \mathcal{L}^\pi(\mathfrak{F})$ and $G/K_2 \in \mathcal{L}^\pi(\mathfrak{F})$. Then $G_\pi K_1/K_1 \subseteq FK_1/K_1$ and $G_\pi K_2/K_2 \subseteq FK_2/K_2$, where $G_\pi K_1/K_1$ is a Hall π -subgroup of G/K_1

and $G_\pi K_2/K_2$ is a Hall π -subgroup of G/K_2 , FK_1/K_1 is an \mathfrak{F} -projector of G/K_1 and FK_2/K_2 is an \mathfrak{F} -projector of G/K_2 .

Therefore $G_\pi K_1 \subseteq FK_1$ and $G_\pi K_2 \subseteq FK_2$. Hence $G_\pi K_1 \cap G_\pi K_2 \subseteq FK_1 \cap FK_2$. By [18, Lemma 1.4] and [10, Theorem 15.2] we have $G_\pi(K_1 \cap K_2) \subseteq F(K_1 \cap K_2)$, i.e., $G_\pi \subseteq F$. Thus $G \in \mathcal{L}^\pi(\mathfrak{F})$. This proves (1).

(2) Inclusion $\mathcal{L}^\pi(\mathfrak{F}) \subseteq \mathfrak{E}_{\pi'}\mathcal{L}^\pi(\mathfrak{F})$ is obvious. We show that $\mathfrak{E}_{\pi'}\mathcal{L}^\pi(\mathfrak{F}) \subseteq \mathcal{L}^\pi(\mathfrak{F})$. Let $G \in \mathfrak{E}_{\pi'}\mathcal{L}^\pi(\mathfrak{F})$. Then $G^{\mathcal{L}^\pi(\mathfrak{F})} \in \mathfrak{E}_{\pi'}$ and $G/G^{\mathcal{L}^\pi(\mathfrak{F})} \in \mathcal{L}^\pi(\mathfrak{F})$.

Let G_π be a Hall π -subgroup of G and let F be an \mathfrak{F} -projector of G . By [10, Lemma 15.2] and [10, Lemma 15.1], we see, $G_\pi G^{\mathcal{L}^\pi(\mathfrak{F})}/G^{\mathcal{L}^\pi(\mathfrak{F})}$ is a Hall π -subgroup of $G/G^{\mathcal{L}^\pi(\mathfrak{F})}$ and $FG^{\mathcal{L}^\pi(\mathfrak{F})}/G^{\mathcal{L}^\pi(\mathfrak{F})}$ is an \mathfrak{F} -projector of $G/G^{\mathcal{L}^\pi(\mathfrak{F})}$. Therefore

$$G_\pi G^{\mathcal{L}^\pi(\mathfrak{F})}/G^{\mathcal{L}^\pi(\mathfrak{F})} \subseteq F^x G^{\mathcal{L}^\pi(\mathfrak{F})}/G^{\mathcal{L}^\pi(\mathfrak{F})}.$$

By [10, Lemma 15.1], $F^x G^{\mathcal{L}^\pi(\mathfrak{F})}/G^{\mathcal{L}^\pi(\mathfrak{F})}$ is an \mathfrak{F} -projector of $G/G^{\mathcal{L}^\pi(\mathfrak{F})}$, where $x \in G/G^{\mathcal{L}^\pi(\mathfrak{F})}$. Consequently,

$$|G/G^{\mathcal{L}^\pi(\mathfrak{F})} : F^x G^{\mathcal{L}^\pi(\mathfrak{F})}/G^{\mathcal{L}^\pi(\mathfrak{F})}| = \frac{|G|}{|F^x G^{\mathcal{L}^\pi(\mathfrak{F})}|} =$$

$$\frac{|G||F \cap G^{\mathcal{L}^\pi(\mathfrak{F})}|}{|F||G^{\mathcal{L}^\pi(\mathfrak{F})}|} = \frac{|G|}{|F||G^{\mathcal{L}^\pi(\mathfrak{F})}|}$$

is a π' -number. Since $|G^{\mathcal{L}^\pi(\mathfrak{F})}|$ is a π' -number, $|G : F|$ is a π' -number. Thus a Hall π -subgroup G_π of G is contained in the \mathfrak{F} -projector F of G . Hence $G \in \mathcal{L}^\pi(\mathfrak{F})$. The lemma is proved. \square

The following theorem shows that if \mathfrak{F} is a saturated formation, then the formation $\mathcal{L}^\pi(\mathfrak{F})$ is saturated.

Theorem. *Let $\mathfrak{F} = LF(F)$ be the formation of all soluble π -groups. Then $\mathcal{L}^\pi(\mathfrak{F}) = LF(f)$ for a local satellite f such that*

$$f(p) = \begin{cases} (\mathfrak{F} \downarrow F(p)), & \text{if } p \in \pi, \\ \mathfrak{S}^\pi, & \text{if } p \in \pi'. \end{cases}$$

Proof. If $\pi = \emptyset$, then $\mathcal{L}^\emptyset(\mathfrak{F}) = \mathfrak{S}^\pi$; if $\pi = \mathbb{P}$, then $\mathcal{L}^\mathbb{P}(\mathfrak{F}) = \mathfrak{F}$. We have saturated formations \mathfrak{S}^π and \mathfrak{F} , and hence the result.

Now suppose $\emptyset \subset \pi \subset \mathbb{P}$. Since a formation \mathfrak{F} is saturated, by [3, IV, 4.3] we have $\text{Char}(\mathfrak{F}) = \sigma(\mathfrak{F})$.

So a π -soluble group G is $\sigma(\mathfrak{F})$ -soluble. Consequently, the subgroup $G^{\mathfrak{F}}$ of G is $\sigma(\mathfrak{F})$ -soluble.

By Lemma 1 we have $\mathfrak{F} = LF(m)$, where m is a local satellite of \mathfrak{F} such that $m(p) = \mathfrak{F} \downarrow F(p)$ for all $p \in \mathbb{P}$.

We show $LF(f) \subseteq \mathcal{L}^\pi(\mathfrak{F})$. Suppose $LF(f) \not\subseteq \mathcal{L}^\pi(\mathfrak{F})$. Let G be a group of minimal order in $LF(f) \setminus \mathcal{L}^\pi(\mathfrak{F})$. Then G is a monolithic group and $K = G^{\mathcal{L}^\pi(\mathfrak{F})}$ is the socle of G . We have $|G/K| < |G|$, so by induction, $G/K \in \mathcal{L}^\pi(\mathfrak{F})$. If T is an \mathfrak{F} -projector of G and G_π is a Hall π -subgroup of G , then by the definition $\mathcal{L}^\pi(\mathfrak{F})$, we have $G_\pi K/K \subseteq TK/K$. Hence $G_\pi K \subseteq TK$. Since G is π -soluble, K is either a p -group, where $p \in \pi$ or a normal π' -subgroup.

Let K be a p -group, where $p \in \pi$. Since $G \in LF(f)$,

$$G/C_G(K) \in f(p) = (\mathfrak{F} \downarrow F(p)).$$

By Lemma 1, an \mathfrak{F} -projector T covers K , i.e., $K \subseteq T$. Therefore $T = TK \supseteq G_\pi K \supseteq G_\pi$. It follows that $G \in \mathcal{L}^\pi(\mathfrak{F})$, a contradiction.

Now let $K \in \mathfrak{E}_{\pi'}$. Lemma 3 implies $G \in \mathfrak{E}_{\pi'} \mathcal{L}^\pi(\mathfrak{F}) = \mathcal{L}^\pi(\mathfrak{F})$, a contradiction.

We prove the converse inclusion, i.e., $\mathcal{L}^\pi(\mathfrak{F}) \subseteq LF(f)$. Suppose $\mathcal{L}^\pi(\mathfrak{F}) \not\subseteq LF(f)$. Let H be a group of minimal order in $\mathcal{L}^\pi(\mathfrak{F}) \setminus LF(f)$. Then H is a monolithic group and $R = H^{LF(f)}$ is the socle of H . Since H is π -soluble, R is either a p -group, where $p \in \pi$ or a normal π' -subgroup.

Let R be a π' -subgroup. By induction, $H/R \in LF(f)$. Consequently, all factors of the chief series $H \supset \dots \supset R$ are f -central. By assumption, $H/C_H(R) \in \mathfrak{S}^\pi = f(p)$. Hence $H \in LF(f)$, a contradiction.

Now let R be a p -group, where $p \in \pi$. If H_π is a Hall π -subgroup of H and V is an \mathfrak{F} -projector of H , then by Chunihin's Theorem [19], we have $R \subseteq H_\pi$. Since $H \in \mathcal{L}^\pi(\mathfrak{F})$, $H_\pi \subseteq V$. Consequently, $R \subseteq V$, i.e., V covers R . Lemma 1 implies that R is m -central chief factor of H . By induction, $H/R \in LF(f)$. Consequently, $H \in LF(f)$. This final contradiction completes the proof. \square

References

- [1] P. Lockett, On the theory of Fitting classes of finite soluble groups, *Math. Z.* **131** (1973) pp. 103–115.
- [2] P. Hauck, Eine Bemerkung zur kleinsten normalen Fittingklasse, *J. Algebra* **53** (1978) pp. 395–401.
- [3] K. Doerk and T. Hawkes, *Finite soluble groups*, Walter de Gruyter, Berlin-New York, 1992.
- [4] O.J. Brison, Hall operators for Fitting classes, *Arch. Math.[Basel]* **80** (33) (1979/80) pp. 1–9.
- [5] E. Cusack, Strong containment of Fitting classes, *J. Algebra* **64** (3) (1980) pp. 414–429.
- [6] O.J. Brison, A criterion for the Hall-closure of Fitting classes, *Bull. Austral. Math. Soc.* **23** (1981) pp. 361–365.

- [7] O.J. Brison, Hall-closure and products of Fitting classes, *J. Austral. Math. Soc. Ser. A.* **32** (1984) pp. 145–164.
- [8] K. Doerk and T. Hawkes, Ein Beispiel aus der Theorie der Schunckklassen, *Arch. Math. [Basel]* **31** (1978) pp. 539–544.
- [9] D. Blessenohl, Über Formationen und Halluntergruppen endlicher auflösbarer Gruppen, *Math. Z.* **142** (3) (1975) pp. 299–300.
- [10] L.A. Shemetkov, *Formations of finite groups*, Nauka, Main Editorial Board for Physical and Mathematical Literature, Moscow, 1978 (in Russian).
- [11] L.M. Slepova, On formations of E^δ -groups, *Doklady AN BSSR* **21** (7) (1977) pp. 587–589 (in Russian).
- [12] E.P. Vdovin, D.O. Revin and L.A. Shemetkov, Formations of finite \mathcal{C}_π -groups, The International Scientific Conference "X The Belarusian Mathematical Conference" November 3–7, 2008, Minsk, Institute of Mathematics of National Academy of Sciences of Belarus (2008) pp. 12–13.
- [13] L.A. Shemetkov and A.F. Vasil'ev, Nonlocal formations of finite groups, *Doklady AN Belarusi* **39** (4) (1995) pp. 5–8 (in Russian).
- [14] Guo Wenbin and Li Baojun, On Shemetkov problem for Fitting classes, *Beitr. Algebra und Geom.* **48** (1) (2007) pp. 281–289.
- [15] N.T. Vorob'ev and V.N. Zagurskii, Fitting classes with given properties of Hall subgroups, *Matematicheskiye Zametki* **78** (2) (2005) pp. 234–240 (in Russian); translated in *Mathematical Notes* **78** (2) (2005) pp. 213–218.
- [16] B. Huppert, *Endliche Gruppen*, Springer–Verlag, Berlin–Heidelberg–New York, 1967.
- [17] A.N. Skiba and L.A. Shemetkov, Multiply ω -local formations and Fitting classes of finite groups, *Matematicheskiye Trudy* **2** (1) (1999) pp. 114–147 (in Russian); translated in *Siberian Adv. Math.* **10** (2) (2000) pp. 112–141.
- [18] N.T. Vorob'ev, Maximal screens of local formations, *Algebra i Logika* **18** (2) (1979) pp. 137–161 (in Russian).
- [19] S.A. Chunihin, *Subgroups of finite groups*, Nauka i tekhnika, Minsk, 1964 (in Russian).

CONTACT INFORMATION

- A. P. Mekhovich** Polotsk State Agricultural Economic College, Oktyabrskaya street, 55, Polotsk, 211413, Belarus
E-Mail: amekhovich@yandex.ru
- N. N. Vorob'ev,**
N. T. Vorob'ev Vitebsk State University of P.M. Masherov, Moscow Avenue, 33, Vitebsk, 210038, Belarus
E-Mail: vornic2001@yahoo.com

Received by the editors: 08.11.2009
and in final form 21.05.2010.